microzone®

MegaMix

P. Code	Reactions (20µl)	Volume	Component	Description	Lot Number	Expiry
2MMB-1	>50	1 ml	MegaMix Blue	1.1X Concentrated, Standard Taq, 220μM dNTPs, 2.75 mM MgCl ₂ and blue loading dye in optimised buffer.		

Applications

- Routine PCR up to 6kb
- Genotyping.
- TA cloning.
- Microarray Analysis.
- Colony PCR.
- Amplification of GC and AT rich templates.
- Methylated DNA.

Product Description

Containing all the components needed to perform PCR swiftly and reliably. The 1.1X mix contains standard Taq DNA polymerase, 220 μ M dNTP, 2.75 mM MgCl₂ and blue agarose loading dye in Microzone's proprietary enhancing buffer. MegaMix Blue is optimised to provide high yields under standard conditions. The blue agarose loading dye incorporated allows easy visualisation and eliminates the need for additional gel loading buffers. The dye does not inhibit restriction enzymes or ligases and does not fluoresce at the wavelengths used by automated DNA sequencers so downstream processes are not impacted.

Prepare a master mix as described in the table below. This reaction can be scaled

Key Features

- More confidence in amplification and PCR test.
- Mastermix format is ready to use.
- Inert blue agarose loading dye allows for easy visualisation.
- Easy set up and PCR optimisation.
- Broad range of templates and conditions.
- Extremely stable—can be freeze thawed many times.



Thermocycling

Transfer the reactions to the thermal cycler and set as follows:

Cycles	Temperature	Time
1	95°C	3 min
25-30	95°C	30 sec
	55-65°C	30 to 60 sec
	72°C	45 to 60 sec

Annealing temperature (55-65°C) may require optimisation depending on the specific primers in use.

The run time can be shortened by optimising the steps of the thermocycling profile.

For research use only

Product Handling

Protocol

Components

MegaMix Blue

Primers

Template

This products is to be used as follows.

Thaw all reagents completely and mix well before use.

according to the quantity of reactions required. Mix gently, avoiding bubbles, centrifuge is necessary. Include a no template control and positive control as required.

Storage

To ensure the quality of the product until the expiry date keep at the recommended storage temperature and limit exposure to light.

Volume

18µl

0.5-1 μl

0.5-1 μl

Contamination Control

To prevent erroneous results ensure work environment is free of contamination by cleaning your workstation and equipment daily with a DNA decontaminant daily, wear gloves, use sterile tubes and filter pipet tips.

Simple | Effective | Efficient